



Quest of novel GH20 N-acetyl hexosaminidasetransglycosylating catalysts: functional screening, data mining and semi-rational mutagenesis

Teze, David; Visnapuu, Triinu; Kjeldsen, Christian; Duus, Jens Øllgaard; Cola, Corinna Schiano di; Lie, Aleksander; Pedersen, Lars Haastrup; Stougaard, Peter; Svensson, Birte

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Teze, D., Visnapuu, T., Kjeldsen, C., Duus, J. Ø., Cola, C. S. D., Lie, A., Pedersen, L. H., Stougaard, P., & Svensson, B. (2017). *Quest of novel GH20 N-acetyl hexosaminidasetransglycosylating catalysts: functional screening, data mining and semi-rational mutagenesis*. Abstract from 19th European Carbohydrate Symposium, Barcelona, Spain.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



QUEST OF NOVEL GH20 *N*-ACETYL HEXOSAMINIDASE TRANSGLYCOSYLATING CATALYSTS: FUNCTIONAL SCREENING, DATA MINING AND SEMI-RATIONAL MUTAGENESIS

David Teze¹, Triinu Visnapuu¹, Christian Kjeldsen², Jens Øllgaard Duus², Corinna Schiano di Cola¹, Aleksander Lie³, Lars Haastrup Pedersen³, Peter Stougaard⁴, Birte Svensson¹

¹ Department of Biotechnology and Biomedicine. Technical University of Denmark. Kongens Lyngby. Denmark.

² Department of Chemistry. Technical University of Denmark. Kongens Lyngby. Denmark.

³ Department of Chemistry and Bioscience. Aalborg University. Denmark.

⁴ Department of Plant and Environmental Sciences. University of Copenhagen. Denmark.
e-mail: david.teze@gmail.com

Lack of access to certain types of oligosaccharides is a severe bottleneck for advances in glycosciences. The transglycosylation activity of retaining glycoside hydrolases (GH) has been used to provide oligosaccharides. The main drawbacks of those enzymes are the competing hydrolysis reaction and the fact that the products are also substrates, thus needing a kinetic control of the reaction. Several approaches have been developed to overcome these, including mechanism modifications (e.g. glycosynthases, chemical rescue), functional screening and data mining to find natural transglycosidases, directed evolution and targeted mutagenesis [1].

Here we focused on *N*-acetyl hexosaminidases from family GH20 that catalyse removal or addition of GlcNAc and GalNAc. Despite sharing a substrate-assisted mechanism with GH85, for which several glycosynthases have been created [2], no successful GH20 glycosynthase has been reported. Thus, we turned to discovery and characterization of new GH20s and performing a systematic mutagenesis study. Several new GH20s of bacterial origin were isolated and described by functional screening and data mining, including transglycosidases able to synthesize lacto-*N*-triose, a valuable oligosaccharide [3], as well as genuine hydrolases. Mutational analysis of all residues within the catalytic domain which were unchanged in >99% of 371 aligned GH20 sequences was pursued. Indeed, it has been shown that targeting conserved residues increases the likelihood of finding advantageous mutations [4,5]. Furthermore, it allows for transfer of successful mutations to other GH20s to find new efficient transglycosidases.

Notably, even though most conserved residues occur within the first and second shell of substrate interaction, 9 residues inside the (β/α)₈ barrel pointing toward the active site are also conserved. To the best of our knowledge, such residues were not studied, although one of them mutated by directed evolution of a GH29 enzyme improved the transglycosylation yield and was transferable to other GH29 members [6]. Here transglycosylation yields of mutants in first shell, second shell and other residues within the (β/α)₈ barrel will be compared for GH20.

Acknowledgments: Innovation Fond Denmark / Danish Council for Strategic Research supported this work.

[1] Slámová K., et al., Synthesis of derivatized chitoooligomers using transglycosidases engineered from the fungal GH20 β -N-acetylhexosaminidase. *Adv. Synth. Catal.* **2015**, 357, 1941–50.

[2] Danby, P.M. and Withers, S.G. Advances in enzymatic glycoside synthesis. *ACS Chem. Biol.* **2016**, 11, 1784–94.

[3] Nyffenegger C., et al. Backbone structures in human milk oligosaccharides: trans-glycosylation by metagenomic β -N-acetylhexosaminidases. *Appl Microbial Biotechnol* **2015**, 99, 7997-8009.

[4] Teze D., et al. Semi-rational approach for converting a GH1 β -glycosidase into a β -transglycosidase. *Protein Eng. Des. Sel.* **2014**, 27, 13–9.

[5] Teze D., et al. Semi-rational approach for converting a GH36 α -glycosidase into an α -transglycosidase. *Glycobiology* **2015**, 25, 420–7.

[6] Saumonneau A., et al. Design of an α -L-transfucosidase for the synthesis of fucosylated HMOs. *Glycobiology* **2016**, 26, 261–9.